

عنوان مقاله:

Persistent sub-diaphragmatic activity on the myocardial perfusion scan with 99mTc-Sestamibi

محل انتشار:

مجله پزشکی هسته ای ایران، دوره 16، شماره 1 (سال: 1387)

تعداد صفحات اصل مقاله: 5

نویسندگان:

Ramin Sadeghi - *Department of Nuclear Medicine, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran*

Vahid Reza Dabbagh Kakhki - *Department of Nuclear Medicine, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran*

Rasoul Zakavi - *Department of Nuclear Medicine, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran*

Mehdi Momennezhad - *Department of Nuclear Medicine, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran*

خلاصه مقاله:

We present a female patient with atypical chest pain who was referred to our department for ischemia evaluation. 99mTc-MIBI myocardial perfusion scan with dipyridamole stress was performed. Sub-diaphragmatic activity in the hepatic tissue and then in the bowel loops caused severe overlap on the inferior wall even on consecutive delayed images. Dipyridamole stress was repeated for the patient with 201Tl. The study was interpretable this time without any interfering sub-diaphragmatic activity.

کلمات کلیدی:

99mTc sestamibi, Myocardial perfusion scan, Sub-diaphragmatic activity

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1008391>

