

عنوان مقاله:

The survey of nauplius and cyprid larvae of *Amphibalanus improvisus* (Darwin, 1854) in different areas of the southern Caspian Sea

محل انتشار:

چهارمین کنگره بین المللی تحقیقات شیلات و آبزیان (سال: 1399)

تعداد صفحات اصل مقاله: 1

نویسندگان:

M Rowshan Tabari - *Caspian Sea Ecology Research Center (CSERC), Sari, Iran*

R Rahmati - *Caspian Sea Ecology Research Center (CSERC), Sari, Iran*

N Khodaparast - *Caspian Sea Ecology Research Center (CSERC), Sari, Iran*

F.S Tahami - *Caspian Sea Ecology Research Center (CSERC), Sari, Iran*

خلاصه مقاله:

This study carried out by the R/V Guilan cruises in the Iranian coast of the Caspian Sea in 2010. We investigated nauplius and cyprid of the barnacle *Amphibalanus improvisus* in the southern Caspian Sea. The aim of this study was to identify and determine density of Total nauplius and larvae of *A. improvisus* (meroplankton) in the different seasons and regions of the Caspian Sea. The highest abundance of *A. improvisus* was observed in the spring 1347 ± 1254 ind. m^{-3} . The peak of abundance was observed in the spring, declining sharply in the summer and autumn, and rising again in the winter. Annual changes showed that 66% of the abundance of *A. improvisus* existed in the west, 19% in the center and 15% in the east regions. In spring, the highest abundance of nauplius and cyprid larvae was observed. nauplius (82%) and larvae (48%) included population of *A. improvisus*. The purpose of this study was to monitor of *A. improvisus* in Southern Caspian Sea and its abundance was shown in different regions and seasons

کلمات کلیدی:

Zooplankton, *Amphibalanus improvisus*, Nauplius, cyprid, Meroplankton, Caspian Sea

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1134921>

