

عنوان مقاله:

?RRM1 and RRM2 of human dead end protein: which one does interact with p27 transcript

محل انتشار:

نهمین همایش بیوانفورماتیک ایران (سال: 1398)

تعداد صفحات اصل مقاله: 1

نویسندگان:

Homa Faraji - Biosensor Research Center, Endocrinology and Metabolism Molecular-Cellular Sciences Institute,
Tehran University of Medical Sciences, Tehran, Iran

Farnaz Banakar - Biosensor Research Center, Endocrinology and Metabolism Molecular-Cellular Sciences Institute,
Tehran University of Medical Sciences, Tehran, Iran

Azadeh Ebrahim-Habibi - Endocrinology and Metabolism Research Center, Endocrinology and Metabolism Clinical
Sciences Institute, Tehran University of Medical Sciences

خلاصه مقاله:

Dead End (Dnd) is an essential RNA binding protein for maintaining viable germ cells in vertebrates. Binding of Dnd to the 3'-UTRs of some mRNAs such as p27 and LATS2 blocks the inhibitory function of microRNAs from these transcripts and in this way helps to maintain p27 and LATS2 protein expression [1]. Human Dnd has two RNA recognition motifs (RRM) including RRM1 and RRM2. p27 has two Dnd interacting sequences (p27-1 and p27-2) [2]. The aim of this study is modeling and simulating of interactions between human Dnd and p27

کلمات کلیدی:

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1164307>

