

عنوان مقاله:

The Relationship between Binocular Summation in Contrast Sensitivity and Stereopsis: Cross Sectional Study

محل انتشار:

مجله پژوهش در علوم توانبخشی, دوره 17, شماره 1 (سال: 1400)

تعداد صفحات اصل مقاله: 7

نویسندگان: منیرہ محجوب

منیرہ محجوب فرخندہ شھری

خلاصه مقاله:

Introduction: Binocular summation occurs in the presence of normal visual function of both eyes. Stereopsis is considered to be the finest function of binocular vision. Therefore, the aim of this study was to investigate the effect of binocular summation on contrast sensitivity and its relationship with stereopsis. Materials and Methods: In this cross-sectional study, $\mathcal{F} \circ$ students of Zahedan University of Medical Sciences, Zahedan, Iran (IV men and FM women) with a mean age of Y1.Y $\bullet \pm$ 1.F $\circ \bullet$ years were selected by convenience sampling method. Monocular and binocular contrast sensitivity were measured with the best corrected vision in photopic conditions using the Metrovision test for spatial frequencies of \circ .Y \circ , 1.Y \circ , M, \mathcal{F} , F, IM, and Y \circ cycles per degree. Stereopsis was also measured by TNO and Titmus tests. Statistical analysis was performed using the repeated measures analysis of variance (ANOVA) and multiple linear regression. Results: The results of repeated measures ANOVA showed that binocular vision compared to monocular vision significantly increased contrast sensitivity (P < \circ . \circ \circ 1). But there was no significant difference between the contrast sensitivity of right and left eyes (P = \circ .Y $\mathcal{F}\mathcal{F}$). Pearson corralation test did not show a significant relationship between stereopsis measured by crossed and uncrossed methods in Titmus and TNO tests with binocular contrast sensitivity at all spatial frequencies (P = \circ .IIF). Conclusion: Binocular summation in presence of normal binocular function can improve visual functions such as the increased binocular contrast sensitivity compared to monocular function can improve visual functions such as the increased binocular contrast sensitivity compared to monocular .contrast sensitivity which confirms the importance of treating binocular anomalies such as amblyopia

کلمات کلیدی:

Contrast Sensitivity, Binocular vision, Monocular vision, stereopsis, Regression Analysis

لینک ثابت مقاله در پایگاه سیویلیکا:



https://civilica.com/doc/1592692