

## عنوان مقاله:

The Common Trunk of the Celiac and Superior Mesenteric as Celiacomesenteric Trunk: A Case Report

## محل انتشار:

مجله تروما, دوره 27, شماره 5 (سال: 1401)

تعداد صفحات اصل مقاله: 5

## نویسندگان:

.Mohamad Mosahar Mehrnahad - School of Medicine, Yazd University of Medical Sciences, Yazd, Iran

.Mersad Mehrnahad - Department of Radiology, Qom University of Medical Sciences, Qom, Iran

## خلاصه مقاله:

Background: The superior mesenteric and celiac arteries separately originate from the aorta. Developmental changes can lead to the formation of a united celicaomesenteric trunk. Case presentation: An ۸۳-year-old male with epigastric shooting pain radiating to the back was admitted to the hospital. Computed tomography angiography of the abdomen was performed to rule out dissection. Incidentally, a united celiacomesenteric trunk was observed. A long right renal artery directly originated from the aorta with a corkscrew appearance, and the left suprarenal and the left gastric arteries arose from the aorta with a common trunk. Conclusion: A united celiacomesenteric trunk can put the individual at risk of surgical complications and ischemia due to the lack of dual supply

## کلمات کلیدی:

celiacomesenteric trunk, Abdominal aorta, Anatomic variation, abdominal vasculature

## لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1704251>

