

## عنوان مقاله:

Risk Factors Affecting Anxiety and Depression in Malignancy Patients in Makassar

## محل انتشار:

مجله علوم دارویی و شیمی، دوره 7، شماره 1 (سال: 1403)

تعداد صفحات اصل مقاله: 10

## نویسندگان:

Tutik Harjianti - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Sitti Rahmah - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Andi Fachruddin Benyamin - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Syakib Bakri - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Hasyim Kasim - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Wasis Udaya - *Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Saidah Syamsuddin - *Department of Psychiatry, Faculty of Medicine, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

Arifin Seweng - *Department of Biostatistics, Faculty of Public Health, Hasanuddin University, Makassar, ۹۰۲۴۵, Indonesia*

## خلاصه مقاله:

**Objective:** This study aimed to determine the risk factors that affect anxiety and depression in patients diagnosed with malignancy in Makassar. **Materials and methods:** A cross-sectional study was conducted involving ۵۵ patients with malignancy who met the research criteria. Assessment of anxiety and depression was conducted using the GAD-۷ and PHQ-۹ instruments. The statistical test employed the Chi-Square test. The results were considered significant if the p-value < ۰.۰۵. **Results:** The study found that ۴۹.۱% of the patients had anxiety, ۶۱.۸% had depression, and ۴۵.۵% had a combination of anxiety and depression. The analysis showed a significant relationship between performance status and anxiety ( $p < ۰.۰۱$ ), performance status and depression ( $p < ۰.۰۵$ ), as well as performance status with mixed anxiety and depression ( $p < ۰.۰۰۱$ ). **Conclusion:** The findings of this study suggest that performance status ECOG  $\geq 3$  is a risk factor for anxiety, depression, and both anxiety and depression in patients diagnosed with malignancy.

**کلمات کلیدی:**

anxiety, depression, Haematological malignancy, Malignancy, solid tumour

**لینک ثابت مقاله در پایگاه سیویلیکا:**

<https://civilica.com/doc/1822832>

