

## عنوان مقاله:

Disseminated fusariosis with ecthyma gangrenosum-like lesions in a refractory acute myeloid leukemia patient

## محل انتشار:

سرطان معده، دوره 5، شماره 1 (سال: 1398)

تعداد صفحات اصل مقاله: 5

## نویسندگان:

Masoud Mardani - *Infectious diseases and tropical medicine research center , Shahid Beheshti university of medical sciences, Tehran, Iran*

Rozita Khodashahi - *Infectious diseases and tropical medicine research center , Shahid Beheshti university of medical sciences, Tehran, Iran*

Ensieh Lotfali - *Department of Medical Parasitology and Mycology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Sara Abolghasemi - *Infectious diseases and tropical medicine research center , Shahid Beheshti university of medical sciences , Tehran, Iran*

Mojdeh Hakemi-Vala - *Department of microbiology, Medical school, Shahid Beheshti university of medical sciences , Tehran, Iran*

## خلاصه مقاله:

**Background and Purpose:** Fusarium species is an opportunistic mold that causes disseminated infections in immunocompromised patients. Given the high mortality rate of this infection, it is important to make a definite diagnosis when encountering suspected cases. **Case report:** Herein, we presented a ۳۵-year-old man diagnosed with acute myeloid leukemia with a prolonged febrile neutropenic period and ecthyma gangrenosum-like lesions, along with fungemia and disseminated fusariosis. The patient died despite receiving combination therapy, perhaps due to the nonrecovery of neutrophil. **Conclusion:** Ecthyma gangrenosum-like lesions due to disseminated fusariosis might be easily misdiagnosed. Consequently, more attention should be paid to the cutaneous lesions in immunocompromised patients.

## کلمات کلیدی:

Disseminated fusariosis, Ecthyma gangrenosum-like lesions, Fungemia

## لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1838111>



