# عنوان مقاله:

Prevalence of helicobacter pylori in patients undergoing upper GI endoscopy in hospital No \ Kerman university of medical sciences

## محل انتشار:

مجله دانشگاه علوم پزشكى كرمان, دوره 1, شماره 3 (سال: 1373)

تعداد صفحات اصل مقاله: 5

#### نویسنده:

H Jojaji - Assistant professor

### خلاصه مقاله:

Prevalence of helicobacter pylori was evaluated by urease test in ۵\mathbb{N} patients undergoing GI endoscopy, in hospital No\ Kerman university of medical sciences and health services from June \99\mathbb{N} through April \99\mathbb{N}. H. Pylori was detected in \partial \89\mathbb{N} of tested samples by urease test. There was no significant correlation with sex, age and cigarette smoking. Prevalence of H.pylori was \89\% in control group, \\000c0\% for esophageal varices, \mathbb{V}\cdot\% for gastric ulcer, \mathbb{Y}\cdot\% for prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for non-ulcer dyspepsia, \\partial\nabla\) for duodenitis, \(\mathbb{P}\cdot\nabla\) for prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for non-ulcer dyspepsia, \\partial\nabla\) for duodenitis, \(\mathbb{P}\cdot\nabla\) for prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for and duodenal ulcer groups compared to control was higher (statistically insignificant). Prevalence of H.pylori in females suffering from duodenal ulcer was also higher than males (\gamma\cdot\nabla\) for compared to \(\mathbb{V}\cdot\nabla\) for prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for medical and the prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for model and ulcer was also the prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for the prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for model and ulcer was also the prepyloric ulcer, \(\mathbb{P}\cdot\nabla\) for the prepyloric ulcer, \(\mathbb{

## كلمات كليدي:

Helicobacter, duodenal ulcer, peptic ulcer, Gastritis, duodenitis

لینک ثابت مقاله در پایگاه سیویلیکا:

https://civilica.com/doc/1935294

