

عنوان مقاله:

The Study of Self-mutilation in Patients with Psychiatric Disorders Admitted to the Lavasani Hospital in ۲۰۱۴-۲۰۱۵

محل انتشار:

مجله قانون طب, دوره 1, شماره 1 (سال: 1398)

تعداد صفحات اصل مقاله: 5

نویسندگان:

Medical Student Research Committee, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran - - -

Department of Psychiatry, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran - - -

Student Research Committee, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran - - -

خلاصه مقاله:

Introduction: Self-mutilation can be defined as a deliberate behavior that directly damages the body and damages body tissues or even amputations which, for a few minutes, deprives a person of the feeling or misery position. This study was performed to examine different types of self-mutilation in patients suffering from psychiatric disorders and its relation with the type of illness among patients attending to Lavasani Hospital in ۲۰۱۴-۱۵. Methods: This study is a cross-sectional observation. Forty-two patients suffering from psychiatric illnesses who had performed self-mutilation entered the study. The prevalence of the type of self-mutilation, the used device, and its relation with the type of mental disorder was analyzed. Results: The Results showed that ۹۰.۵% of patients used sharp objects, ۴.۸% used fire, ۲.۴% used rocks, and ۲.۴% used other devices for self-mutilation. ۷۶.۲% of patients injured the organs, ۱۹% injured the head and neck, ۲۴% injured the body, and ۲.۴% injured the stomach area. Age and sex did not have a direct correlation with the place and device used ($P > ۰.۰۵$), but type of disorder did show a significant relationship with the place and device used ($P = ۰.۰۰۱$). Conclusion: It is concluded that self-mutilation in mental patients is done mainly using sharp objects in the organs.

کلمات کلیدی:

Psychiatric Diseases, self-mutilation, neurosis

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1979955>

