

عنوان مقاله:

Evaluation of suicide attempts with drug poisoning in North-West of Iran

محل انتشار:

مجله اورژانس و تروما, دوره 1, شماره 1 (سال: 1394)

تعداد صفحات اصل مقاله: 2

نویسندگان:

Samad Shams Vahdati - Department of Emergency Medicine, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Narges Moradi - School of Nursing, Tabriz University of Medical Sciences, Tabriz, Iran

Jamil Hemat Ghadim - Department of Law, Payame Noor University of Tabriz, Tabriz, Iran

Shahrad Tajoddini - Kerman Neuroscience Research Center, Kerman University of Medical Sciences, Kerman, Iran

خلاصه مقاله:

Deliberate Self-Poisoning (DSP) or drug induced suicide is the most common method of suicide in the developed countries and is associated with significant morbidity and mortality (۱-۳). Factors associated with DSP include economic and social factors, culture, religion, and health (۴-۶). Other (symptoms that may influence suicide encompass demographic information like age, sex, and, ethnicity (۶-۸)

کلمات کلیدی:

Suicide, Drug poisoning, Iran

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/2038535>

