سیویلیکا – ناشر تخصصی مقالات کنفرانس ها و ژورنال ها گواهی ثبت مقاله در سیویلیکا CIVILICA.com

عنوان مقاله:

The frequency of using different types of pacifier and bottle nipple among \-\Y\ months old children

محل انتشار:

مجله بهداشت دهان و دندان و اپیدمیولوژی دهان و دندان, دوره 5, شماره 1 (سال: 1395)

تعداد صفحات اصل مقاله: 4

نویسندگان:

Hamidreza Poureslami - Professor, Oral and Dental Diseases Research Center AND Kerman Social Determinants on Oral Health Research Center AND
,Department of Pediatric Dentistry, School of Dentistry, Kerman University of Medical Sciences, Kerman

Razieh Hoseinifar - Assistant Professor, Department of Operative Dentistry, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

Reihaneh Hosseinifar - Resident, Department of Pediatric Dentistry, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

Hamid Sharifi - Modeling in Health Research Center, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

Parnian Poureslami - Student of Dentistry, School of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran

خلاصه مقاله:

BACKGROUND AND AIM: Sucking of some types of a pacifier may be a risk factor for mal-development of orofacial structures and malocclusion. This study assessed the prevalence of using of different types of pacifier and bottle nipple among \-\text{1-14} months old children in Kerman, Iran.METHODS: In this cross-sectional study, a total of \(\text{r··}\text{ monthers interviewed at r· specialized private pediatric offices in Kerman. The mothers had \-\text{1-14} months old children.} A checklist includes items about pacifier sucking, bottle feeding habits, as well as mothers' ability to recognize different kinds of available pacifier and bottle nipple was used.RESULTS: The rate of pacifier-sucking was \(\text{rv.r/\%}\), and use of bottle feeding was \(\text{rv.r/\%}\), and among of the users, \(\text{rf.·}\) and \(\text{rd.r/\%}\) were used orthodontic (functional) types of pacifier and bottle nipple respectively. However, only \(\text{rd.r/\%}\) of mothers had adequate ability to recognize orthodontic pacifier and bottle nipple were low, and a few mothers had adequate ability to recognize their differences

كلمات كليدى:

Pacifier, Bottle Feeding, Bottle Nipple, Breast feeding, Sucking Habits

لینک ثابت مقاله در پایگاه سیویلیکا:

https://civilica.com/doc/2046456

