

عنوان مقاله:

Evaluating the quality of life in patients with ulcerative oral lesions

محل انتشار:

مجله بهداشت دهان و دندان و اپیدمیولوژی دهان و دندان، دوره 8، شماره 4 (سال: 1398)

تعداد صفحات اصل مقاله: 6

نویسندگان:

Fatemeh Lavaee – Assistant Professor, Department of Oral and Maxillofacial Medicine, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran

Azita Sadeghzadeh – MSc Student, Department of Oral and Maxillofacial Medicine, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran

Bahar Afrooz – Department of Oral and Maxillofacial Medicine, School of Dentistry, Yasuj University of Medical Sciences, Yasuj, Iran

Ali Golkari – Assistant Professor, Department of Oral Health and Social Dentistry, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran

Abdollah Piri-Zarrini – Student Research Committee, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran

خلاصه مقاله:

**BACKGROUND AND AIM:** Oral mucosal lesions can affect patient's quality of life (QOL). In this evaluation, Persian version of Chronic Oral Mucosal Disease Questionnaire (COMDQ) was used to assess participants' QOL. **METHODS:** This cross-sectional study was done during ۲۰۱۵-۲۰۱۶ in School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran. ۹۵ participants with pemphigus vulgaris (PV), oral recurrent aphthous stomatitis (RAS), and oral lichen planus (OLP) enrolled in this study. The Persian version of COMDQ with ۲۶ questions and ۴ domains was used for assessing QOL. The data were analyzed in SPSS software. P-value less than ۰.۰۵ was considered significant. Independent t-test, analysis of variance (ANOVA), and Tukey's test were used to assess scores of QOL. **RESULTS:** The final QOL scores were  $۴۵.۹۵ \pm ۱۶.۳۱$ ,  $۵۲.۲۸ \pm ۱۷.۶۴$ , and  $۵۰.۰۲ \pm ۱۷.۳۶$  for men, women, and all patients, respectively. Patients with OLP and RAS had good QOL, but patients with PV reported lower level of QOL (moderate). None of the COMDQ domains showed significant correlation with gender except pain and functional limitation and overall QOL score. **CONCLUSION:** The result of this evaluation revealed a good QOL; considering the type of oral disease, QOL ranged between moderate for patients with PV and good for patients with OLP and RAS.

کلمات کلیدی:

Lichen Planus, stomatitis, Aphthous, pemphigus, Quality of Life

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/2048393>

