

عنوان مقاله:

Generalization of Katugampola fractional kinetic equation involving incomplete H-function

محل انتشار:

مجله روشهای محاسباتی برای معادلات دیفرانسیل، دوره 12، شماره 4 (سال: 1403)

تعداد صفحات اصل مقاله: 15

نویسندگان:

Nishant - - Department of Mathematics, Malaviya National Institute of Technology Jaipur, India

Sanjay Bhattar - Department of Mathematics, Malaviya National Institute of Technology Jaipur, India

Sunil Purohit - Department of HEAS (Mathematics), Rajasthan Technical University, India. Department of Computer Science and Mathematics, Lebanese American University, Beirut, Lebanon

خلاصه مقاله:

In this article, Katugampola fractional kinetic equation (KE) has been expressed in terms of polynomials along with incomplete H-function, incomplete Meijer's G-function, incomplete Fox-Wright function, and incomplete generalized hypergeometric function, weighing the novel significance of the fractional KE that appear in a variety of scientific and engineering scenarios. T-Laplace transform is used to solve the Kathugampola fractional KE. The obtained solutions have been presented with some real values and the simulation was done via MATLAB. Furthermore, the numerical and graphical interpretations are also mentioned to illustrate the main results. Each of the obtained conclusions is of a general nature and is capable of generating the solutions to several fractional KE

کلمات کلیدی:

Fractional kinetic equation, Incomplete H-functions, Mellin-Barnes type contour

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/2088608>

