

عنوان مقاله:

Molecular typing of Leishmania major parasite in the main vector of leishmaniasis in Khuzestan by targeting the ITSrDNA gene

محل انتشار:

هشتمین همایش بیوتکنولوژی جمهوری اسلامی ایران و چهارمین همایش ملی امنیت زیستی (سال: 1392)

تعداد صفحات اصل مقاله: 2

نویسندگان:

.Roozbeh Taslimian - Molecular Systematics Laboratory, Deparment of Parasitology, Pasteur Institute of Iran, Tehran

.Parviz Parvizi - Molecular Systematics Laboratory, Deparment of Parasitology, Pasteur Institute of Iran, Tehran

Dr. Ahmad Reza Esmaili - Molecular Systematics Laboratory, Deparment of Parasitology, Pasteur Institute of Iran, .Tehran

Adel Spotin - Molecular Systematics Laboratory, Deparment of Parasitology, Pasteur Institute of Iran, Tehran

خلاصه مقاله:

Background & Objective:Leishmania major is an underlying agent of zoonotic cutaneous leishmaniasis in an endemic focus of Khuzestan. In this study, we diagnose Leishmania parasites by targeting of ITS-rDNA gene because there is no molecular typing and characterization which has been carried out on P. papatasi as a main vector of L. major yetMaterials & Methods:Sandflies were caught using sticky traps and light traps. After dissecting, sandflies wereidentified. DNA was extracted and ITS-rDNA gene of Leishmania in sandflies amplified byPCR methods and by RFLP and sequencing then using molecular software Leishmania majorwas definitively approved.Results:Among 1500 sandflies, 198 female P. papatasi were selected as gravid, semi gravid and freshfed. DNA was extracted from all of them and the ITS-rDNA gene was amplified. About 8percent of sandflies was infected with Leishmania parasites. After RFLP and sequencing samples Leishmania major was identified in all cases.Conclusions:The sequences results .obtained in this study compared with similar cases recorded in the GenBank and confirmed

كلمات كليدي:

Leishmania major, Sandfly, Khuzestan, ITS-rDNA, PCR-RFLP

لینک ثابت مقاله در پایگاه سیویلیکا:

https://civilica.com/doc/377122

