

عنوان مقاله:

The Role of Gender in Cholecystitis Complications

محل انتشار:

مجله بین المللی تحقیقات پیشرفته زیست شناختی و زیست پزشکی، دوره 2، شماره 6 (سال: 1393)

تعداد صفحات اصل مقاله: 4

نویسندگان:

Nozar Dorostan - *Department of surgery, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran*

Mohammad Reza Boostani - *Department of Radiology, Golestan Hospital, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran*

Iraj Nazari - *Department of surgery, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran*

Elham Rajai - *Department of Internal Medicine, Golestan Hospital, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran*

خلاصه مقاله:

Cholecystitis presents as acute or chronic. Severity of cholecystitis depends on several factors. Aim of this study was to evaluate severity of laparoscopic cholecystectomy findings based on gender type. In a retrospective and epidemiological study, a total of 101 patients who were underwent laparoscopic cholecystectomy were selected from 2001 to 2006 in university affiliated hospitals of Ahwaz, Iran. Among 101 patients were assessed in our study, 41(40.6%) were men and 60(59.4%) were women. Evaluating of patients according to sex revealed that proportion of empyema, gallbladder stones, gangrene, switch to open cholecystectomy, and acute cholecystitis were different statistically and were higher in men than women. But chronic cholecystitis was significantly lower in men than women. As our findings demonstrated, we suggest that men with cholecystitis need rapid surgical intervention to prevent and reduce complications of disease

کلمات کلیدی:

Gender, Cholecystitis, Laparoscopy

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/443073>

