

عنوان مقاله:

Exsanguinating Hemorrhage during Open Biopsy in a Primary Breast Angiosarcoma: A Case Report

محل انتشار:

دوازدهمین کنگره بین المللی سرطان پستان (سال: 1394)

تعداد صفحات اصل مقاله: 1

نوپسندگان:

Majid Akrami - MD Breast Diseases Research Center, Department of Surgical Oncology, Shiraz University of Medical .Sciences, Shiraz, Iran

Mastoureh Mohammadipour - General surgeon, Ziaei hospital, Ardakan, Yazd

.Maral Mokhtari - MD Department of radiotherapy, Kermanshah university of medical sciences, Kermanshah,Iran

.Malihe Dayani - MD Department of Pathology, Nemazee Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

خلاصه مقاله:

Introduction: Angiosarcomas are endothelial cell neoplasms in the lining of the blood vessel wall and account for about 0.04% of all breast malignancies with a high rate of error in primary diagnosis. The breast angiosarcoma is a rare and uncommon pathology and hasbeen described mostly as case reports. Indeed, only a limited number of cases have been published. Accordingly, the natural history of this tumor and its clinical course remain unclear, and as a consequence, no uniform treatment strategy exists. Methods: We present the clinical course and challenges in the diagnosis of a primary angiosarcoma of the breast in a young woman, presenting with a mass in her left breast. Fineneedle aspiration and core needle biopsy failed to confer a correct diagnosis. Results: She suffered severe bleeding at the time of open biopsy and underwent total mastectomy, followed by adjuvant chemotherapy Conclusion: Young women with solid breast tumors, especially those that are highly vascular, should be considered malignant until proven otherwise. Accurate diagnosis may be difficult. Open biopsy can be diagnostic, although exsanguinating bleeding may occur. Thus, performing open biopsy in locations other than equippedoperating rooms may be hazardous and should be avoided

کلمات کلیدی:

لینک ثابت مقاله در پایگاه سیویلیکا:

https://civilica.com/doc/740732

