

## عنوان مقاله:

Convulsion Associated with Gastroenteritis

## محل انتشار:

مجله پزشکی بالینی، دوره 5، شماره 3 (سال: 1397)

تعداد صفحات اصل مقاله: 3

## نویسندگان:

Maryam Khalesi - *Department of Pediatrics, Ghaem Medical Center, Mashhad University of Medical Sciences, Mashhad, Iran*

Farhad Heydarian - *Department of Pediatrics, Ghaem Medical Center, Mashhad University of Medical Sciences, Mashhad, Iran*

Seyed Javad Sayedi - *Department of Pediatrics, Ghaem Medical Center, Mashhad University of Medical Sciences, Mashhad, Iran*

Shima Badzai - *Department of Pediatrics, Ghaem Medical Center, Mashhad University of Medical Sciences, Mashhad, Iran*

## خلاصه مقاله:

Convulsion with mild gastroenteritis is an afebrile seizure associated with viral gastroenteritis in a healthy child without fever, dehydration, electrolyte imbalance, meningitis, or encephalitis. Convulsion with mild gastroenteritis is more common in children aged 1 to 2 years. Usually, Convulsions are brief generalized tonic clonic type. Most convulsions occur within first 24 hours of illness onset. Rotaviral gastroenteritis is known as the most common type of gastroenteritis associated with Convulsion. Laboratory investigations are normal. Also EEG and neuroimaging are usually normal. Long term antiepileptic treatment is not necessary. It is usually a benign condition with good prognosis and no risk for developing epilepsy in future. Considering this etiology of seizure could prevent supernumerary evaluations and long-term antiepileptic treatment.

## کلمات کلیدی:

Child, Convulsion, Gastroenteritis

## لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/892431>

