

عنوان مقاله:

Association Between Vitamin D Level in Pregnant Women and Level of TSH in Their Neonates

محل انتشار:

پانزدهمین کنگره بین المللی زنان و مامایی ایران (سال: 1398)

تعداد صفحات اصل مقاله: 1

نویسنده:

Shahnaz Ahmadi - Iran university of medical science . Firoozabadi hospital

خلاصه مقاله:

Background: prevention of hypovitaminosis D IN pregnancy and related fetal and maternal adverse effects would help to improve the outcomes. However in this study, the association between vitamin D level in pregnant women and level of TSH in their neonates was determined. Material and Methods: In this observational study as comparative cross-sectional, 100 consecutive pregnant women from firoozabadi hospital were enrolled and the serum vitamin D was assessed in them and the association with neonatal TSH was determined. Results: in this study the mean maternal serum vitamin D level was 24.9 ±13.3 and the mean neonatal serum TSH level was 2.8 ±1. There was no statistically significant association between vitamin D level in pregnant women and TSH level in their neonates (p> 0.05). Conclusion: It may be concluded that Vitamin D level in pregnant women and level of TSH in their neonates are not interrelated

کلمات کلیدی: vitamin D , Thyroid ,neonates

لینک ثابت مقاله در پایگاه سیویلیکا:

https://civilica.com/doc/954074

