

## عنوان مقاله:

The Importance of Screening in Prevention of Postpartum Depression

## محل انتشار:

پانزدهمین کنگره بین المللی زنان و مامایی ایران (سال: 1398)

تعداد صفحات اصل مقاله: 1

## نویسندگان:

Mohammad Arzanlou - M.D., Psychiatrist, Khoy University of medical science, Khoy Iran

Zoleykha Asgarlou - MSc of Midwifery, Department of Midwifery, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Zanjan, Iran

## خلاصه مقاله:

Dear Editor-in-Chief: Postpartum Depression (PPD) is a major public health problem and a common clinical disorder that occurs after childbirth (1). This disorder is characterized by symptoms such as depressed mood, disappointment, irritability, anxiety, lack of enjoyment of life, excessive fatigue, sleep and appetite disorders, mental and motor disturbances, disruption of family and social communication, self-harming and feeling of guilty or inadequacy particularly regarding the infant care (2-4). In addition to the problems for the mother, the baby and other family members are also affected by this problem. Psychological and social factors are considered as main factors of PPD (5). As a result, these two factors should be focused in order to resolve the consequent problems. Screening is one of methods through which some of the causing factors can be identified and by preventing their occurrence, the incidence of this type of depression can be reduced and more patients can be found and treated as well. In other words, a wide range of the health system, from primary health centers to providers and therapists, plays an important role in this regard. Generally, there are tools for PPD screening, and many countries use screening, however these screenings are usually not mandatory (6). Given that depression is increasing in recent years and the World Health Organization has ranked the major depression as the fourth leading cause of disease burden among all diseases, which can lead to life with disability (7), different countries should more seriously seek the prevention and implement screening. Especially in low and middle-income countries that in which less services are usually provided to people and those with a poor economic status are less looking for programs such as screening. Furthermore, in most countries, poor economic status, for various reasons, leads to a low level of people's awareness of health-related programs. Culture and traditions are also other important factors. In some countries, especially in the less developed regions, there is no regular referral culture for prenatal care and referral to a physician. In screening, risk factors of PPD should be noted. The risk factors include previous history of PPD, neurological disorders before childbirth, stressful signs of depression during pregnancy and childbirth, low socioeconomic status, poor social support, problems such as pre-eclampsia, chronic diseases of the mother and embryonic issues (8). Given that screening and ... treatment of postpartum depression are cost-effective interventions and should be

## کلمات کلیدی:

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/954085>



